

## Administering Vaccines to Adults: Dose, Route, Site, Needle Size, and Preparation

Vaccine	Dose	Route	Site	Needle Size	Vaccine Preparation
<b>Tetanus, Diphtheria (Td) with Pertussis (Tdap)</b>	0.5 mL	IM	Deltoid	22–25g, 1–1½"	Shake vial vigorously to obtain a uniform suspension prior to withdrawing each dose. Whenever solution and container permit, inspect vaccine visually for particulate matter and/or discoloration prior to administration. If problems are noted (e.g., vaccine cannot be resuspended), the vaccine should not be administered.
<b>Hepatitis A (HepA)</b>	≤18 yrs.: 0.5 mL ≥19 yrs.: 1.0 mL	IM	Deltoid	22–25g, 1–1½"	
<b>Hepatitis B (HepB)</b>	≤19 yrs.: 0.5 mL ≥20 yrs.: 1.0 mL	IM	Deltoid	22–25g, 1–1½"	
<b>HepA+HepB (Twinrix)</b>	≥18 yrs.: 1.0 mL	IM	Deltoid	22–25g, 1–1½"	
<b>Human papillomavirus (HPV)</b>	0.5 mL	IM	Deltoid	22–25g, 1–1½"	
<b>Influenza, trivalent inactivated (TIV)</b>	0.5 mL	IM	Deltoid	22–25g, 1–1½"	
<b>Pneumococcal polysaccharide (PPV)</b>	0.5 mL	IM	Deltoid	22–25g, 1–1½"	
		SC	Fatty tissue over triceps	23–25g, 5/8"	
<b>Meningococcal, conjugated (MCV4)</b>	0.5 mL	IM	Deltoid	22–25g, 1–1½"	<p>Reconstitute just before using. Use only the diluent supplied with the vaccine. Inject the volume of the diluent shown on the diluent label into the vial of lyophilized vaccine and gently agitate to mix thoroughly. Withdraw the entire contents and administer immediately after reconstitution.</p> <p>Discard single dose MPSV4, varicella, and zoster vaccines if not used within 30 minutes after reconstitution.</p> <p><b>Note:</b> Unused reconstituted MMR vaccine and multidose MPSV4 vaccine may be stored at 35–46°F (2–8°C) for a limited time. The reconstituted MPSV4 vaccine must be used within 35 days; the reconstituted MMR vaccine must be used within 8 hours. Do not freeze either reconstituted vaccine.</p>
<b>Meningococcal, polysaccharide (MPSV4)</b>	0.5 mL	SC	Fatty tissue over triceps	23–25g, 5/8"	
<b>Measles, mumps, rubella (MMR)</b>	0.5 mL	SC	Fatty tissue over triceps	23–25g, 5/8"	
<b>Zoster (Zos)</b>	0.65 mL	SC	Fatty tissue over triceps	23–25g, 5/8"	
<b>Varicella (Var)</b>	0.5 mL	SC	Fatty tissue over triceps	23–25g, 5/8"	
<b>Influenza, live, attenuated (LAIV)</b>	0.5 mL (0.25 mL into each nostril)	Intranasal spray	Intranasal	NA	

**Please note:** Always refer to the package insert included with each biologic for complete vaccine administration information. CDC’s Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well. Access the ACIP recommendations at [www.immunize.org/acip](http://www.immunize.org/acip).